



COCKTAIL MELASMA - 1,8% TRANEXAMIC ACID-

DESCRIPTION

0/

This cocktail is recommended for hyperpigmented skin caused by melasma. Tranexamic acid is capable of inhibiting hiperpigmentation by inhibiting various enzimatic pathways.

On the one hand it competes with Tyrosine, having more affinity than Tyrosine for the enzyme Tyrosinase. In this way the melanin formation pathway is blocked.

On the other hand, it blocks the formation of plasmin, a protein that induces the synthesis of the hormone that activates melanogenesis (Alpha-melano-stimulant) and the release of arachidonic acid that would end up activating melanogenesis.

FUNITIONS

| % | ACTIVE INGREDIENTS | FUNTIONS |
|-------------------|--------------------|--|
| 1,8 | TRANEXAMIC ACID | Depigmentation capacity. Inhibits tyrosinase enzyme regulating the formation of melanin pigment. |
| | | |
| AVAILABLE FORMATS | | |
| | 8 | |
| OBSERVATIONS | | SHELF LIFE AND STORAGE |

A OTIVE INCORPORATE

Topaz packaging Expiry date: 3 Years from production

PAO: 3 months

Storage: Storage between 15-25°C. Protect from sunlight.

Sterilization:

The information contained in this document is prepared for informational and commercial purposes only, referring to the product described. The information regarding raw materials and active ingredients, may vary for reasons beyond our Company's control.

Do not use this information for: regulatory purposes, preparation of safety dossier, product registration or printing of labels and packaging. For these purposes, please contact the Commercial Department of Neftis Laboratorios SL.

Ver: 06 Pàgina 1 de 1 Neftislaboratorios.com